



Clubroot Factsheet

Series 1 for farms

sheet 4

Managing New and Isolated Outbreaks

Fact: ISOLATED OR SMALL SPOT OUTBREAKS OF CLUBROOT CAN BE ERADICATED

Dealing with an isolated or spot outbreak of clubroot:

- **Identify** the disease.
- **Contact** local departments of Primary Industries or if required use diagnostic services (eg. Crop Health Services ph: (03) 92109222) for some assistance.



Typical spot outbreak of clubroot



- **Quarantine** by marking off an area at least 3 metres from the outer edge of the infected plants.
- **Remove** all plants within this area.
- **Dispose** by digging up diseased plants and bagging the roots and attached soil.
- **Discard** diseased plants by burning or disposing in an industrial landfill.

Fungicides will not kill established disease.

The information contained in this factsheet resulted from research conducted under the NATIONAL CLUBROOT PROJECT. Supported by DPI (VIC), Agriculture WA, DPIWE (TAS), DPI (QLD) and NSW Agriculture.

Disclaimer: This publication may be of assistance to you but the State of Victoria and its officers do not guarantee that the publication is without flaw of any kind or is wholly appropriate for your particular purposes and therefore disclaims all liability for any error, loss or other consequence which may arise from you relying on any information in this publication.

© State of Victoria, Department of Primary Industries, June 2005.

This publication is copyright. No part may be reproduced by any process except in accordance with the provisions of the Copyright Act 1968.